

## Reply to comment by L. Fenoglio-Marc et al. on “On the steric and mass-induced contributions to the annual sea level variations in the Mediterranean Sea”

David García,<sup>1</sup> Benjamin F. Chao,<sup>2</sup> Jorge Del Río,<sup>3</sup> Isabel Vigo,<sup>1</sup>  
and Jesús García-Lafuente<sup>3</sup>

Received 26 July 2007; accepted 24 September 2007; published 27 December 2007.

**Citation:** García, D., B. F. Chao, J. Del Río, I. Vigo, and J. García-Lafuente (2007), Reply to comment by L. Fenoglio-Marc et al. on “On the steric and mass-induced contributions to the annual sea level variations in the Mediterranean Sea,” *J. Geophys. Res.*, **112**, C12019, doi:10.1029/2007JC004465.

[1] *García et al.* [2006] (hereafter referred to as G06) study the water mass variations in the Mediterranean Sea using a direct and an indirect method. The direct method is based on the gravitational signature of the monthly water mass variations estimated by the space mission of GRACE. The indirect method is as follows: The sea level variation (SLV) is a result of water mass change (SLVmass) plus density variations (SLVsteric). Therefore the residual signal between the SLV (estimated by altimetry measurements) and the SLVsteric (estimated by temperature and salinity profiles from the ECCO ocean model) yields the SLVmass. G06 found a good agreement between the results of the two methods.

[2] However, subsequent similar studies, notably *Fenoglio-Marc et al.* [2006] (hereafter referred to as F06) found different results, which are commented on by *Fenoglio-Marc et al.* [2007] (hereafter referred to as F07). F07 found some differences between G06 and F06 in both SLVsteric and SLVmass signals.

[3] In response to F07, we make three points.

[4] 1. F07 states that the SLVsteric signal in G06 is an overestimate. We thus made a revision of the program used to estimate that signal and did identify a wrong line in our code. Once it was corrected, we obtained a similar signal to that in F06. We thank F07 for pointing out the discrepancy.


[5] 2. Both G06 and F06 use the same monthly GRACE data, namely the Release 01 (RL01) of level-2 CSR GRACE gravity fields between April 2002 and July 2004, and apply the same treatment to the data in relation to the nontidal barotropic ocean correction, and degree-1 and degree-2 spherical harmonics. However, there are three main differences in estimating the average SLVmass over the Mediterranean: (1) the radius and the truncation degree used in the averaging Gaussian filter [*Swenson and Wahr*, 2002]; (2) the spectral leakage in SLVmass produced by

water mass variations in the Mediterranean surrounding areas; and (3) restoration of the diminution of the SLVmass signal produced by the Gaussian filter. We address them in inverse order, as follows.

[6] The Gaussian filter diminishes the amplitude of the averaged signal. F06 restored that diminution according to the algorithm of *Velicogna and Wahr* [2006], which unfortunately was not yet published when G06 was accepted for publication.

[7] Spectral leakage is a main theoretical extension of F06 over G06, and obviously a justified improvement.

[8] High-degree SH coefficients from GRACE are contaminated by noise [e.g., *Tapley et al.*, 2004; *Wahr et al.*, 2004; *Swenson and Wahr*, 2006], and then, an averaging Gaussian filter is used to reduce their contribution [*Swenson and Wahr*, 2002]. The averaging Gaussian filter depends on a radius,  $r$ . G06 uses  $r = 1000$  km and SH coefficients up to degree 15, and F06 uses  $r = 400$  km and SH coefficients up to degree 90. F07 states that filter in F06 is more appropriated than the one in G06. However, we do not agree with this statement. Figure 1 shows that both filters produce almost




**Figure 1.** Average Gaussian filtered GRACE SLVmass over the Mediterranean Sea using two radii,  $r = 400$  km (red line) and  $r = 1000$  km (blue line).


<sup>1</sup>Space Geodesy Laboratory, Applied Mathematics Department, University of Alicante, Alicante, Spain.

<sup>2</sup>College of Earth Sciences, National Central University, Taiwan, China.


<sup>3</sup>Department of Applied Physics II, University of Malaga, Malaga, Spain.



**Figure 2.** Percentage of the accumulative amplitude spectrum, as a function of degree, of the averaging Gaussian filters in the Mediterranean Sea using  $r = 1000$  km (blue line) and  $r = 400$  km (red line).



**Figure 3.** Weight of the SH coefficients when a Gaussian filter is applied for two radii:  $r = 400$  km (red line) and  $r = 1000$  km (blue line).



**Figure 4.** Annual amplitude of the GRACE-deduced mass variations. SH coefficients have been Gaussian filtered with two radii: (a)  $r = 1000$  km and (b)  $r = 400$  km.

exactly the same signal. The reason is found in Figure 2, which shows the percentage of the accumulative amplitude spectrum, as a function of degree, of the averaging Gaussian filters in the Mediterranean Sea using  $r = 1000$  km (blue line) and  $r = 400$  km (red line). The truncation at degree 15 in G06 accounts for the 95% of the signal and then it is not relevant. In the case of F06 the use of SH coefficients of degree further 50 is completely useless.

[9] The election of the radius seems not to be very important when applying an averaging Gaussian filter in the Mediterranean Sea. However, when a spatial averaging is not concerned, a Gaussian filtering with  $r = 1000$  km is much more convenient than with  $r = 400$  km. The Gaussian filter weights the SH coefficients as shown in Figure 3, for

$r = 400$  km (red line) and  $r = 1000$  km (blue line). RL01 CSR GRACE SH coefficients are sufficiently well determined up to degree  $\sim 15$  [Tapley *et al.*, 2004; Wahr *et al.*, 2004]. When  $r = 1000$  km is used, SH coefficients between degree 15 and 25 have a very small contribution, and those of degree greater than 25 are virtually zero. However, SH coefficients of degree  $> 15$ , which are already within the range of the noisy SH coefficients, contribute a lot when  $r = 400$  km is used. Figures 4a and 4b show the annual amplitude of GRACE-deduced mass variations using radius of 1000 km and 400 km, respectively, to Gaussian filter the SH coefficients. The resulting map in Figure 4a shows recognizable signals as those of Amazon basin or the Monsoons. Figure 4b shows unrealistic north-south stripes,

which have been related to errors in high-degree SH coefficients [Swenson and Wahr, 2006].

[10] 3. The monthly water mass flux in the Mediterranean can be obtained by simply subtracting the water mass of two consecutive months from the GRACE data. However, GRACE data used in G06 and F07 had gaps between April 2002 and July 2004, in particular, June 2002, July 2002, December 2002, January 2003 and June 2003 are missing. Therefore only 18 monthly flux can be recovered. On the other hand, although F06 states that 20 monthly GRACE data are used, their Figures 3 and 4 show 25 monthly data. We do not know the origin of those extra points and cannot comment on the differences.

[11] **Acknowledgments.** We acknowledge Alberto Escapa for discussions. This work is supported by the Spanish Science and Technology Ministry Project ESP2006-11357.

## References

- Fenoglio-Marc, L., J. Kusche, and M. Becker (2006), Mass variation in the Mediterranean Sea from GRACE and its validation by altimetry, steric and hydrologic fields, *Geophys. Res. Lett.*, *33*, L19606, doi:10.1029/2006GL026851.
- Fenoglio-Marc, L., J. Kusche, M. Becker, and I. Fukumori (2007), Comment on "On the steric and mass-induced contributions to the annual sea level variations in the Mediterranean Sea," *J. Geophys. Res.*, *112*, C12018, doi:10.1029/2007JC004196.
- García, D., B. F. Chao, J. Del Río, I. Vigo, and J. García-Lafuente (2006), On the steric and mass-induced contributions to the annual sea level variations in the Mediterranean Sea, *J. Geophys. Res.*, *111*, C09030, doi:10.1029/2005JC002956.
- Swenson, S., and J. Wahr (2002), Methods for inferring regional surface mass anomalies from Gravity Recovery and Climate Experiment (GRACE) measurements of time-variable gravity, *J. Geophys. Res.*, *107*(B9), 2193, doi:10.1029/2001JB000576.
- Swenson, S., and J. Wahr (2006), Post-processing removal of correlated errors in GRACE data, *Geophys. Res. Lett.*, *33*, L08402, doi:10.1029/2005GL025285.
- Tapley, D. B., S. Bettadpur, M. Watkins, and C. Reigber (2004), The gravity recovery and climate experiment: Mission overview and early results, *Geophys. Res. Lett.*, *31*, L09607, doi:10.1029/2004GL019920.
- Velicogna, I., and J. Wahr (2006), Measurements of time-variable gravity show mass loss in Antarctica, *Science*, *311*, 1754–1756, doi:10.1126/science.1123785.
- Wahr, J., S. Swenson, V. Zlotnicki, and I. Velicogna (2004), Time-variable gravity from GRACE: First results, *Geophys. Res. Lett.*, *31*, L11501, doi:10.1029/2004GL019779.
- B. F. Chao, College of Earth Sciences, National Central University, Cheng-li, Taiwan.
- J. Del Río and J. García-Lafuente, Department of Applied Physics II, University of Malaga, E-29071 Malaga, Spain.
- D. García and I. Vigo, Space Geodesy Laboratory, Applied Mathematics Department, University of Alicante, E-03080 Alicante, Spain. (d.garcia@ua.es)